

On Singularity Formation Under Mean Curvature Flow

I.M.Sigal

Toronto

Joint work with Wenbin Kong, Zhou Gang,
and Dan Knopf

Some of it uses ideas from work with
Dimitra Antonopoulou and Georgia Karali

Heraklion

April 2012



Mean Curvature Flow

The mean curvature flow of a hypersurface $M_0 \subset \mathbb{R}^{d+1}$ is a family of hypersurfaces $M_t \subset \mathbb{R}^{d+1}$ whose smooth immersions $\psi(\cdot, t) : N \rightarrow M_t \subset \mathbb{R}^{d+1}$ satisfy the partial differential equation

$$\partial_t \psi(z, t) = -h(\psi(z, t))$$

where $h(y)$ is the mean curvature vector of M_t at a point $y \in M_t$, with the initial condition $\psi(z, 0) = \psi_0(z)$, an immersion of M_0 .

Applications and Connections

- ▶ Material Science (interface motion between different materials or different phases).
- ▶ Image recognition.
- ▶ Connection to the Ricci flow.
- ▶ Topological classification of surfaces and submanifolds.

Some Key Works: Existence

- ▶ First mathematical treatment (using geometric measure theory): Brakke [1978];
- ▶ Short time existence: Brakke, Huisken, Evans and Spruck, Ilmanen, Ecker and Huisken [1991];
- ▶ Evans and Spruck, Chen, Giga and Goto [1991]: Weak solutions;

Some Key Works: Singularities

The most interesting problem here is formation of singularities.

- ▶ Mu-Tao Wang, Kuo-Wei Lee and Yng-Ing Lee [2008]: Higher Co-dimensions;
- ▶ Huisken [1984]: Collapse of convex hypersurfaces;
- ▶ Grayson, Ecker, Huisken, M. Simon, Dziuk and Kawohl, Smoczyk, Altschuler, Angenent and Giga, Soner and Souganidis [1990-1995]: Neckpinching for periodic, rotationally symmetric;
- ▶ Huisken and Sinestrari [2007-2009]: MCF with surgery and topological classification of surfaces and submanifolds;
- ▶ B. White [2000], Colding and Minicozzi [2012]: Nature and Hausdorff dimension of the singular set.

Classification of Singularities

There are three exact solutions of MCF:

- ▶ Collapsing Euclidean spheres with radii decreasing as $\sqrt{2d(t_* - t)}$;
- ▶ Collapsing Euclidean cylinders with radii decreasing as $\sqrt{2(d - 1)(t_* - t)}$;
- ▶ Static planes.

Huisken conjectured that generic singularities are spheres and cylinders.

Colding and Minicozzi [2012] showed that the only 'linearly' stable self-similar solutions (shrinkers) are spheres.

Stability of Shrinking Spheres

Theorem. (W. Kong-I.M.S.) Let a surface M_0 be close to S^d in H^s , $s > \frac{d}{2} + 1$. Then $\exists t_* < \infty$, s.t. MCF has a solution M_t for $0 \leq t < t_*$ and

- ▶ $M_t \rightarrow z_*$, for some z_* , as $t \rightarrow t_*$;
- ▶ M_t are defined by immersions of S^d ,

$$X(\omega, t) = z(t) + R(\omega, t)\omega,$$

for some $z(t) \in \mathbb{R}^{d+1}$ and $R(t) \in H^s(\Omega)$, satisfying

$$z(t) = z_* + O(\tau^\alpha)$$

$$R(t) = \sqrt{2d} \tau \left(1 + O_{H^s}(\tau^\beta) \right),$$

with $\tau := t_* - t$, $\alpha := \frac{1}{2a}(d + \frac{1}{2} - \frac{1}{2d})$ and $\beta := \frac{1}{2a}(1 - \frac{1}{2d})$.

Previous Results

- ▶ Colding and Minicozzi [2012] : 'linear' stability of spherical shrinkers;
- ▶ Antonopoulou - Georgia Karali - IMS: asymptotics of volume preserving MCF near spheres

Graphs over Cylinders

Our next result deals with initial conditions M_0 , which are graphs over d -dimensional cylinders C^d along the x_{d+1} -axis in \mathbb{R}^{d+1} ,

$$X(\omega, x, t) = (u(\omega, x, t)\omega, x).$$

It combines two results, one with Zhou Gang on equivariant graphs (surfaces of revolution), i.e.

$$u(\omega, x, t) \text{ is independent of } \omega,$$

and one in general case with Zhou Gang and Dan Knopf, which is in preparation. For brevity, the initial conditions are formulated informally.

Theorem. (Zhou Gang-I. M. S., Zhou Gang-D. Knopf- IMS) Let $d \geq 2$ and M_0 be a surface close to a cylinder, \mathcal{C}^d ,

M_0 has an arbitrary shallow waist and is even w.r.to the waist.

Then M_t is defined by an immersion

$$X(\omega, x, t) = (u(\omega, x, t)\omega, x)$$

of C^d , where $(\omega, x) \in \mathcal{C}^d$ and $u(\omega, x, t)$ satisfies

- (i) There exists a finite time t^* such that $\inf u(\cdot, t) > 0$ for $t < t^*$ and $\lim_{t \rightarrow t^*} \inf u(\cdot, t) \rightarrow 0$;
- (ii) If $u_0 \partial_x^2 u_0 \geq -1$ then there exists a function $u_*(\omega, x) > 0$ such that $u(\omega, x, t) \geq u_*(\omega, x)$ for $\mathbb{R} \setminus \{0\}$ and $t \leq t^*$.

Dynamics of Scaling Parameter

Theorem. (Zhou Gang-I. M. S., Zhou Gang-D. Knopf- IMS)

(iii) There exist C^1 functions $\zeta(\omega, x, t)$, $\lambda(t)$, $a(t)$ and $b(t)$ such that

$$u(\omega, x, t) = \lambda(t) \left[\left(\frac{2(d-1) + b(t)y^2}{a(t)} \right)^{\frac{1}{2}} + \zeta(\omega, y, t) \right]$$

with $y := x/\lambda(t)$, $\|\langle y \rangle^{-m} \partial_y^n \zeta(\omega, x, t)\|_\infty \leq cb^2(t)$, $m+n=3$,

(iv) The parameters $\lambda(t)$, $a(t)$ and $b(t)$ satisfy the estimates

$$\lambda(t) = (t^* - t)^{\frac{1}{2}}(1 + o(1));$$

$$b(t) = -\frac{d-1}{\ln|t^*-t|} \left(1 + O\left(\frac{1}{|\ln|t^*-t||^{3/4}} \right) \right);$$

$$a(t) = 1 + \frac{1}{\ln|t^*-t|} \left(1 + O\left(\frac{1}{|\ln|t^*-t||} \right) \right).$$

Comparison with Previous Results

A result similar to (ii) (axi-symmetric surfaces) but for a different set of initial conditions was proven by H.M.Soner and P.E.Souganidis.

The previous result closest to ours is that by S. Angenent and D. Knopf on the axi-symmetric neckpinching for the Ricci flow.

Some ideas of the proof are close to those of Bricmont and Kupiainen on NLH.

All works mentioned above deal with *surfaces of revolution* of barbell shapes (*far from cylinders*) which are either compact (Dirichlet b.c.) or periodic (Neumann b.c.).

These works rely on parabolic maximum principle going back to Hamilton and monotonicity formulae for an entropy functional \int_{M_t} backward heat kernel $(x, t)d\mu_t$, due to Huisken and Giga and Kohn.

Key Steps in Proof

Passing to collapse variables;

Collar lemma;

Estimates of the linear evolution;

Bootstrap.

Collapse Variables

We outline key steps in proof in the *axi-symmetric* case. We pass from the original variables x and t to the collapse variables

$$y := \lambda^{-1}(t)(x - x_0(t)) \text{ and } \tau := \int_0^t \lambda^{-2}(s) ds.$$

Important point: we do not fix $\lambda(t)$ and $x_0(t)$ but consider them as free parameters to be found from (MCF).

For a profile $u(x, t)$ we define the new unknown function

$$v(y, \tau) := \lambda^{-1}(t)u(x, t)$$

with $y := \lambda^{-1}(t)x$ and $\tau := \int_0^t \lambda^{-2}(s)ds$. The function v satisfies the equation

$$\partial_\tau v = \left(\frac{1}{1 + (\partial_y v)^2} \partial_y^2 - ay \partial_y + a \right) v - \frac{d-1}{v}, \text{ with } a := -\lambda \partial_t \lambda.$$

Adiabatic Solutions

The rescaled MC equation has the following cylindrical, static (i.e. y and τ -independent) solution

$$V_a := \left(\frac{d-1}{a}\right)^{\frac{1}{2}} \text{ and } a \text{ is constant} \iff u_{cyl}(t).$$

A larger family of approximate solutions: ignore $\partial_\tau v$ and $\partial_y^2 v$ in the equation for $v \implies$

$$ay\partial_y v - av + \frac{d-1}{v} = 0$$

(adiabatic or slowly varying approximation). This equation has the general solution

$$V_{ab} := \left(\frac{2(d-1) + by^2}{2a}\right)^{\frac{1}{2}}$$

with $b \in \mathbb{R}$. In what follows we take $b \geq 0$ so that V_{ab} is smooth. Note that $V_{a0} = V_a$.

Collar Lemma

We introduce the manifold of adiabatic solutions

$$M_{neck} := \{V_{ab} \mid a, b \in \mathbb{R}^+, b \leq \epsilon\}.$$

Lemma

There exist a small neighbourhood \mathcal{U}_{path} of M_{neck} in $C^1([0, T], \langle y \rangle^3 L^\infty)$ and a unique C^1 map $g : \mathcal{U}_{path} \rightarrow C^1([0, \infty), \mathbb{R}) \times C^1([0, \infty), \mathbb{R})$, such that

$$v(y, \tau) = V_{g(v)(\tau)}(y) + \phi(y, \tau),$$

with $g(u)(\tau) = (a(\tau), b(\tau))$, and

$$\phi(\cdot, \tau) \perp 1, \quad a(\tau)y^2 - 1 \text{ in } L^2(\mathbb{R}, e^{-\frac{a(\tau)}{2}y^2} dy).$$

Similarly, for the initial conditions $v_0(y) := \lambda_0^{-1} u_0(\lambda_0 y)$.

The vectors 1 and $(1 - ay^2)$ which are almost tangent vectors to the manifold, M_{neck} , provided b is sufficiently small.

Lyapunov-Schmidt Splitting (Effective Equations)

Substitute $v(y, \tau) = V_{a(\tau), b(\tau)}(y) + \phi(y, \tau)$
into (MCF) to obtain

$$\partial_\tau \phi = -L_{ab} \phi + F_{ab} + N_{ab}(\phi)$$

where L_{ab} is the linear operator given by

$$L_{ab} := -\partial_y^2 + ay\partial_y - 2a + V_{ab}(y)$$

$F_{ab} \approx$ a sum of generators of broken symmetries (the source term)
and $N_{ab}(\phi)$ is a nonlinearity. Remember that

$$\phi(\cdot, \tau) \perp 1, a(\tau)y^2 - 1 \text{ in } L^2(\mathbb{R}, e^{-\frac{a(\tau)}{2}y^2} dy).$$

Project the above equation on $1, a(\tau)y^2 - 1 \implies$ the *equations for the parameters* $a, b \implies$ need to estimate ϕ .

Estimating ϕ

Let $U(\tau, \sigma)$ be the propagator generated by $-L_a$. By Duhamel principle we rewrite the differential equation for $\phi(y, \tau)$ as

$$\phi(\tau) = U(\tau, 0)\phi(0) + \int_0^\tau U(\tau, \sigma)(F + N)(\sigma)d\sigma.$$

Using the *key propagation estimate* ($\tau \geq \sigma \geq 0$)

$$\|\langle z \rangle^{-3} U(\tau, \sigma)g\|_\infty \lesssim e^{-c(\tau-\sigma)} \|\langle z \rangle^{-3}g\|_\infty,$$

where $g \perp 1$, $a(\tau)y^2 - 1$ in $L^2(\mathbb{R}, e^{-\frac{a(\tau)}{2}y^2} dy)$, we obtain

$$M(\tau) \leq M(0) + b^{\frac{1}{2}}(0)P(M(\tau)),$$

where

$$M(\tau) := \max_{\sigma \leq \tau} b^{-\frac{m+n+1}{2}}(\sigma) \|\langle y \rangle^{-m} \partial_y^n \phi(\cdot, \sigma)\|_\infty.$$

Estimating the Linear Propagator. I

Denote the integral kernel of $U(\tau, \sigma)$ by $K(x, y)$. We have the representation

$$K(x, y) = K_0(x, y) \langle e^V \rangle(x, y),$$

where $K_0(x, y)$ is the integral kernel of the operator $e^{-(\tau-\sigma)L_{a0}}$, $L_{a0} := -\partial_y^2 + ay\partial_y - 2a$, $V := L_{ab} - L_{a0}$ and

$$\langle e^V \rangle(x, y) = \int e^{\int_\sigma^\tau V(\omega(s) + \omega_0(s), s) ds} d\mu(\omega).$$

Here $d\mu(\omega)$ is a harmonic oscillator (Ornstein-Uhlenbeck) probability measure on the continuous paths $\omega : [\sigma, \tau] \rightarrow \mathbb{R}$ with the boundary condition $\omega(\sigma) = \omega(\tau) = 0$ and

$$(-\partial_s^2 + a^2)\omega_0 = 0 \text{ with } \omega_0(\sigma) = y \text{ and } \omega_0(\tau) = x.$$

Estimating the Linear Propagator. II

To estimate $U(x, y)$ for $e^{a(\tau-\sigma)} \leq b^{-1/32}(\tau)$ we use the explicit formula

$$K_0(x, y) = 4\pi(1 - e^{-2ar})^{-\frac{1}{2}} \sqrt{a} e^{2ar} e^{-a \frac{(x - e^{-ary})^2}{2(1 - e^{-2ar})}},$$

where $r := \tau - \sigma$, and the bound

$$|\partial_y \langle e^V \rangle(x, y)| \leq b^{\frac{1}{2}} r,$$

which follows from the definition of $\langle e^V \rangle$ and the properties

$$V(y, \tau) \geq 0 \text{ and } |\partial_y V(y, \tau)| \lesssim b^{\frac{1}{2}}(\tau).$$

Then we iterate using the semi-group property \Rightarrow estimate of the remainder ϕ .

Ricci flow;

Anisotropic mean curvature flow
(Euclidean spheres \Rightarrow Wulff shapes);

Kähler surfaces;

Going through the singularity.

Thank-you for your attention.

Comparison with Yang-Mills and Wave Maps Equations

Compare the dynamics for the scaling parameter $\lambda(t)$ for (MCF) and the critical Yang-Mills equation

$$\lambda \ddot{\lambda} = \frac{3}{4} \dot{\lambda}^4,$$

which gives

$$\lambda \approx \sqrt{\frac{2}{3}} \frac{t_* - t}{\sqrt{-\ln(t_* - t)}}.$$

and the critical wave map equation

$$\dot{\lambda}^2 = \lambda \ddot{\lambda} \ln \frac{a}{\lambda \ddot{\lambda}}, \quad a = 0.122.$$

MCF for Surfaces of Revolution

If $d \geq 2$ and M_0 is a surface of revolution around the axis $x = x_{d+1}$, given by a map $r = u_0(x)$ where $r = (\sum_{j=1}^d x_j^2)^{\frac{1}{2}}$, then M_t is also a surface of revolution and, as long as it is smooth, M_t is defined by the map $r = u(x, t)$, satisfying the PDE

$$\begin{aligned}\partial_t u &= \frac{\partial_x^2 u}{1 + (\partial_x u)^2} - \frac{d-1}{u} & \text{(MCF)} \\ u(x, 0) &= u_0(x).\end{aligned}$$

Our goal is to describe the phenomenon of collapse of such surfaces.

We say $u(x, t)$ collapses at time t^* if $\| \frac{1}{u(\cdot, t)} \|_\infty < \infty$ for $t < t^*$ and $\| \frac{1}{u(\cdot, t)} \|_\infty \rightarrow \infty$ as $t \rightarrow t^*$.

Precise restrictions on the initial conditions

$u_0(x)$ is even and satisfy for $(m, n) = (3, 0)$, $(\frac{11}{10}, 0)$, $(1, 2)$ and $(2, 1)$ the estimates

$$\|u_0(x) - (\frac{2(d-1)+\varepsilon_0 x^2}{2\varsigma_0})^{\frac{1}{2}}\|_{m,n} \leq C\varepsilon_0^{\frac{m+n+1}{2}},$$

$$u_0(x) \geq \frac{1}{\sqrt{2\varsigma_0 + \frac{\varepsilon_0}{d-1}}} g(\sqrt{2\varsigma_0 + \frac{\varepsilon_0}{d-1}}x, \frac{\varepsilon_0}{2\varsigma_0}),$$

$\langle x \rangle^{-1} u_0 \in L^\infty$, $\partial_x u_0 \in L^\infty$, $|\partial_x u_0 u_0^{-\frac{1}{2}}| \leq \kappa_0 \varepsilon_0^{\frac{1}{2}}$, $|\partial_x^n u_0| \leq \kappa_0 \varepsilon_0^{n/2}$, $n = 2, 3$
for some C , $\kappa_0 \geq 2$, and $\frac{1}{2} \leq \varsigma_0 \leq 2$.

Comparison with Ricci Flow

A compact surface of revolution can be given by the metric

$$g = \varphi^2 dx^2 + u^2 g_{can} \text{ on } (-1, 1) \times S^d,$$

where g_{can} is the round metric of radius 1 on S^d (a $SO(d+1)$ -invariant metric on S^{d+1}). ($u(x, t) > 0$ may be regarded as the radius of the hypersurface $\{x\} \times S^d$.)

In this representation the Ricci flow for surfaces of revolution is

$$\partial_t u = \partial_s^2 u - (d-1) \frac{1 - (\partial_s u)^2}{u},$$

where $s(x) := \int_0^x \varphi(y) dy$.

(MCF can also be written as a flow of the metric g .)

Non-compact, non-periodic surfaces are allowed;

Initial conditions include in particular surfaces whose mean curvature changes sign and which might have many necks;

The subleading term in the asymptotic is determined and the remainder is estimated.